

## Travel-Related Arboviral Infections: Chikungunya, Dengue, and Zika Viruses

With its emergence in the Western hemisphere, chikungunya virus infection quickly became another significant source of travel-related arboviral infection among Philadelphia residents along with dengue virus, the most common arboviral infection worldwide. The introduction of Zika virus in Brazil during 2015 and its subsequent spread throughout the Americas and Caribbean make it another travel-related arboviral infection of concern. These infections can cause febrile illness in travelers returning from tropical and subtropical regions, with polyarthralgia occurring in most with chikungunya. Dengue infections can progress to more severe illness (hemorrhagic fever or shock syndrome) that can be fatal. While most individuals infected with Zika virus have mild illness or remain asymptomatic, infection in pregnant women may lead to congenital infection that results in neurologic abnormalities or fetal loss.

Chikungunya, dengue, and Zika viruses are spread by *Aedes spp.* mosquitoes, which are daytime biters and found around homes due to their short flight range. *Aedes aegypti*, a principle vector for transmission of these viruses is not found in Philadelphia. *Aedes albopictus* (Asian tiger mosquito) are present and active during warmer months in Philadelphia. PDPH will continue to assess the presence of *A. albopictus* in Philadelphia and closely monitor human surveillance data to promptly identify local transmission should it occur.

### Chikungunya, Dengue and Zika Infections, Philadelphia, 2011—2016

Year, n	Travel-Related Arboviral Infections		
	Chikungunya	Dengue	Zika
2016 <sup>a</sup>	0	0	—
2015 <sup>a</sup>	2	5	—
2014	42	0	—
2013	0	11	—
2012	0	1	—
2011	0	1	—
Median Age (Range), y	42.5 (5–78)	44 (10–79)	—
Female, n (%)	34 (77)	10 (56)	—
Foreign Born, n (%)	31 (70)	5 (28)	—
Hospitalized, n (%)	9 (20)	13 (72)	—
Death, n (%)	0 (0)	1 (6)	—

<sup>a</sup>Preliminary year to date data.

### Worldwide Distribution Reports

[Chikungunya](#) | [Dengue](#) | [Zika](#)

#### Prevention:


- Advise patients traveling to tropical/subtropical regions to take mosquito-bite prevention measures.
- Advise patients at risk for severe disease (pregnant women or women planning to become pregnant, >65 years, persons with underlying conditions) to postpone or cancel travel to areas with ongoing outbreaks.
- Advise cases to stay indoors and avoid mosquitoes for the first 7 days of illness.

#### Treatment:


- Manage chikungunya and Zika as dengue with acetaminophen recommended for initial treatment.
- If dengue has been ruled out, NSAIDs or narcotics may be considered to manage fever and pain.

### Travel Destinations

#### Chikungunya Cases



#### Dengue Cases



#### Testing and Reporting:

- If either chikungunya, dengue, or Zika is suspected, collect serum and test for all 3 pathogens.
  - Order PCR between day 1 and 8 of illness
  - Order IgM & IgG if ≥ day 4 of symptom onset
- Testing is available through the PA Department of Health's Bureau of Laboratories and for chikungunya and dengue, through Focus Diagnostics. Quest Diagnostics and LabCorp will forward orders to Focus.
- Pregnant women who have traveled to areas with ongoing Zika outbreaks should be monitored for Zika infection. See [CDC's testing algorithm](#).
- Report suspected and confirmed infections to PDPH by telephone at 215-685-6740 (215-686-4514 after hours) or fax at 215-238-6947.