

Table 1. Reported Cases of Sexually Transmitted Diseases and Rates per 100,000 Population: Philadelphia, 1990-2020.

Year	Total Syphilis (All Stages)		Primary & Secondary Syphilis		Early Latent Syphilis		Late & Late Latent Syphilis		Congenital Syphilis		Gonorrhea		Chlamydia ³	
	Cases	Rate ¹	Cases	Rate ¹	Cases	Rate ¹	Cases	Rate ¹	Cases	Rate ²	Cases	Rate ¹	Cases	Rate ¹
1990	6,558	412.5	2,361	148.5	3,907	245.8	111	7.0	179	606.6	19,554	1,230.0	-----	-----
1991	5,201	327.2	1,410	88.7	3,323	209.0	167	10.5	301	1,035.5	15,429	970.5	1,151	-----
1992	4,811	302.6	907	57.1	3,348	210.6	285	17.9	271	975.6	11,914	749.4	8,716	548.3
1993	3,752	236.6	515	32.5	2,829	178.4	255	16.1	153	573.5	10,580	667.3	10,053	634.0
1994	2,394	151.0	298	18.8	1,708	107.7	282	17.8	106	408.5	8,026	506.2	9,956	627.9
1995	1,693	106.8	199	12.6	1,100	69.4	329	20.7	65	271.6	6,565	414.0	8,079	509.5
1996	1,298	81.9	141	8.9	839	52.9	255	16.1	63	276.9	6,415	404.6	8,118	512.0
1997	1,091	68.8	108	6.8	648	40.9	300	18.9	35	158.8	6,504	410.2	10,480	660.9
1998	807	50.9	89	5.6	407	25.7	287	18.1	24	108.8	7,271	458.6	11,763	741.9
1999	826	52.0	69	4.3	394	24.8	355	22.3	8	36.9	7,776	489.1	12,660	796.4
2000	622	41.0	67	4.4	265	17.5	281	18.5	9	41.5	8,170	538.4	13,593	895.7
2001	639	42.1	77	5.1	236	15.6	322	21.2	4	19.1	8,061	531.2	13,586	895.3
2002	589	38.8	71	4.7	221	14.6	293	19.3	4	19.1	7,277	479.5	15,234	1,003.8
2003	587	38.7	98	6.5	194	12.8	292	19.2	3	14.3	5,731	377.6	17,747	1,169.5
2004	470	31.0	72	4.7	151	10.0	247	16.3	0	0.0	5,206	343.1	16,723	1,102.0
2005	417	27.5	86	5.7	122	8.0	207	13.6	2	9.2	5,053	333.0	15,577	1,026.5
2006	540	35.6	125	8.2	184	12.1	231	15.2	0	0.0	5,218	343.8	17,199	1,133.3
2007	500	33.0	136	9.0	203	13.4	152	10.0	9	38.2	5,246	345.7	17,029	1,122.1
2008	526	34.7	150	9.9	178	11.7	191	12.6	7	29.7	4,950	326.2	17,012	1,121.0
2009	704	46.4	218	14.4	255	16.8	215	14.2	4	16.9	4,823	317.8	18,104	1,193.0
2010	667	43.7	238	15.6	223	14.6	189	12.4	1	4.2	6,533	428.1	19,428	1,273.1
2011	698	45.7	207	13.6	242	15.9	227	14.9	4	17.1	6,761	443.1	20,471	1,341.5
2012	798	52.3	269	17.6	261	17.1	233	15.3	5	21.7	7,293	477.9	20,803	1,363.2
2013	961	63.0	278	18.2	332	21.8	299	19.6	1	4.3	6,304	413.0	19,570	1,282.2
2014	892	58.5	279	38.8	333	21.8	247	16.2	4	16.0	5,961	390.6	18,935	1,240.8
2015	916	60.0	314	20.6	368	24.1	230	15.1	4	17.4	6,260	410.2	19,169	1,256.1
2016	1,089	71.4	395	54.9	494	32.4	162	10.6	5	20.3	6,967	456.5	19,992	1,310.1
2017	1,256	82.3	459	30.1	567	37.2	224	14.7	6	28.2	7,288	477.6	21,119	1,383.9
2018	1,214	79.6	408	26.7	540	35.4	263	17.2	3	11.7	7,205	472.2	20,206	1,324.1
2019	1,262	82.7	470	30.8	495	32.4	291	19.1	6	24.9	7,043	461.5	20,354	1,333.8
2020	1,374	90.0	511	33.5	596	39.1	261	17.1	6	25.8	7,302	478.5	15,834	1,037.6

¹ Rates for 1990 to 1999 were calculated using 1990 Census data. Rates for 2000 to 2009 were calculated using 2000 Census data. Rates for 2010 forward were calculated using 2010 Census data.

² Rates of congenital syphilis are calculated per 100,000 live births. (Note: Estimated numbers of live births may be utilized for the year of the report and the year prior.)

³ Chlamydia became a reportable condition in Philadelphia as of October 2nd, 1991.