Philadelphia Department of Public Health Division of Disease Control



as probable Zika.

THOMAS A. FARLEY, MD. MPH Health Commissioner

CAROLINE JOHNSON MD Acting Deputy Health Commissioner & Director, Division of Disease Control

## Zika and Other Arbovirus Activity in Philadelphia November 18, 2016

The introduction of Zika virus in Brazil during 2015 and its subsequent spread throughout Central/South America, the Caribbean, and Mexico make it another travel-related arboviral infection of concern along with dengue and chikungunya. While most individuals infected with Zika virus have mild illness or remain asymptomatic, infection in pregnant women may lead to congenital infection that results in neurologic abnormalities or fetal loss.

|                     | Arboviral Infections, 2016 Philadelphia, PA |         |             |  |  |
|---------------------|---|---------|-------------|--|--|
|                     | Zika <sup>a</sup>                           | Dengue  | Chikungunya |  |  |
|                     |   |         |             |  |  |
| Cases <sup>b</sup>  | 36  | 4       | 0           |  |  |
|                     |   | •       | Ũ           |  |  |
| Travel-associated   | 36°   | 4       | 0           |  |  |
| Locally-acquired    | 0   | 0       | 0           |  |  |
| Median Age          | 34  | 46.5    | —           |  |  |
| (Range), y          | (11—73)                                     | (11—65) |             |  |  |
| Female, n (%)       | 26 (72)                                     | 1 (25)  |             |  |  |
| Foreign Born, n (%) | 26 (72)                                     | 1 (25)  |             |  |  |



<sup>a</sup>Zika cases acquired through sexual contact with a <sup>a</sup>Revised Zika case definition implemented September 26, 2016. Unspecified flavivirus infections re-classified returning traveler excluded from figure.

<sup>b</sup>Preliminary year to date data. Includes confirmed and probable cases.

<sup>c</sup>Two cases acquired through sexual contact with a returning traveler. All others traveled to an affected area.

## Worldwide Distribution Reports: Zika | Dengue | Chikungunya

## Aedes Species Mosquito Activity – 2016 Season

Zika, dengue, and chikungunya viruses are spread by Aedes spp. mosquitoes, which are daytime biters and found around homes due to their short flight range. A.aegypti, a principle vector for transmission of these viruses is not found in Philadelphia. A. albopictus (Asian tiger mosquito), a less efficient vector are present and active during warmer months in Philadelphia. PDPH will continue to assess the presence of A. albopictus in Philadelphia and closely monitor A. albopictus Locations<sup>a</sup> human surveillance data to promptly identify local transmission should it occur.

|  | A. aegypti | A. albopictus | Philadelphia, 2016  |
|--|------------|---------------|---|
| Number of identifications since May 2016                       | 0          | 698           |   |
| edian number of mosquitoes per trap (range)                    | —          | 5<br>(1—1138) |   |
| ercentage of all mosquito traps set during 2016 ason           | —          | 41.0%         |   |
| ercentage of residential zip codes with <i>Aedes</i> osquitoes | —          | 97.9%         |   |
| umber of Aedes mosquito pools tested for Zika                  | —          | 217           | 53 48 1   |
| umber of Zika positive Aedes mosquito pools                    | —          | 0             | <sup>a</sup> Adult A. <i>albopictus</i> identified 698 times at 2 |

 Report suspected and confirmed infections to the Philadelphia Department of Public Health by telephone at 215-685-6742 (215-686-4514 after hours) or fax at 215-238-6947.

 For more information on the diagnosis and prevention of Zika and other arboviruses, visit: https://hip.phila.gov/DiseaseControlGuidance/DiseasesConditions/Arboviruses/Zika