



Philadelphia Department of Public Health
Division of Disease Control

THOMAS A. FARLEY, MD, MPH
 Health Commissioner

CAROLINE JOHNSON, MD
 Deputy Health Commissioner

STEVEN ALLES, MD, MS
 Director, Division of Disease Control

Zika and Other Arbovirus Activity in Philadelphia
September 13, 2017

The introduction of Zika virus in Brazil during 2015 and its subsequent spread throughout Central/South America, the Caribbean, and Mexico make it another travel-related arboviral infection of concern along with dengue and chikungunya. While most individuals infected with Zika virus have mild illness or remain asymptomatic, infection in pregnant women may lead to congenital infection that results in neurologic abnormalities or fetal loss.


	Arboviral Infections, Philadelphia, PA, 2016—2017		
	Zika ^a	Dengue	Chikungunya
Cases ^b	56	3	0
2016	48	3	0
2017	8	0	0
Travel-associated	56 ^c	3	0
Locally-acquired	0	0	0
Median Age (Range), y	34 (<1—73)	35 (11—58)	—
Female, n (%)	41 (73)	1 (33)	—
Foreign Born, n (%)	38 (68)	1 (33)	—

^aRevised Zika case definition implemented September 26, 2016. Unspecified flavivirus infections re-classified as probable Zika.

^bPreliminary year to date data. Includes confirmed and probable cases.

^cTwo cases acquired through sexual contact with a returning traveler. Three cases were infants with probable, asymptomatic congenital infection and no identified birth defects. All others traveled to an affected area.

Travel Destinations
**Arboviral Infections, Philadelphia, PA
 2016—2017**




Worldwide Distribution Reports
[Zika](#) | [Dengue](#) | [Chikungunya](#)

***Aedes Species* Mosquito Activity – 2017 Season**

Zika, dengue, and chikungunya viruses are spread by *Aedes spp.* mosquitoes, which are daytime biters and found around homes due to their short flight range. *A. aegypti*, a principle vector for transmission of these viruses is not found in Philadelphia. *A. albopictus* (Asian tiger mosquito), a less efficient vector are present and active during warmer months in Philadelphia. PDPH will continue to assess the presence of *A. albopictus* in Philadelphia and closely monitor human surveillance data to promptly identify local transmission should it occur.

***A. albopictus* Locations^a**



^aAdult *A. albopictus* identified 855 times at 354 locations.

	<i>A. aegypti</i>	<i>A. albopictus</i>
Number of identifications since May 2017	0	855
Median number of mosquitoes per trap (range)	—	6 (1—814)
Percentage of all mosquito traps set during 2017 season	—	60%
Percentage of residential zip codes with <i>Aedes</i> mosquitoes	—	94%

- Report suspected and confirmed infections to the Philadelphia Department of Public Health by telephone at 215-685-6742 (215-686-4514 after hours) or fax at 215-238-6947.
- For more information on the diagnosis and prevention of Zika and other arboviruses, visit: <https://hip.phila.gov/DiseaseControlGuidance/DiseasesConditions/Arboviruses/Zika>