Point of Care Testing Observations (e.g., assisted blood glucose monitoring)									
HH performed		Clean gloves worn		Single use, lancet used? ¹		Testing meter ²		Gloves removed ³	HH performed ³
O Yes O	No	O Yes	O No	O Yes	O No	0 0	Dedicated to resident, cleaned/disinfected before storing Cleaned/disinfected before next resident	O Yes O No	O Yes O No
O Yes O	No	O Yes	O No	O Yes	O No	0 0	Dedicated to resident, cleaned/disinfected before storing Cleaned/disinfected before next resident	O Yes O No	O Yes O No
O Yes O	No	O Yes	O No	O Yes	O No	0 0	Dedicated to resident, cleaned/disinfected before storing Cleaned/disinfected before next resident	O Yes O No	O Yes O No
O Yes O	No	O Yes	O No	O Yes	O No	0 0	Dedicated to resident, cleaned/disinfected before storing Cleaned/disinfected before next resident	O Yes O No	O Yes O No
O Yes O	No	O Yes	О No	O Yes	O No	0 0	Dedicated to resident, cleaned/disinfected before storing Cleaned/disinfected before next resident	O Yes O No	O Yes O No
O Yes O	No	O Yes	O No	O Yes	O No	0 0	Dedicated to resident, cleaned/disinfected before storing Cleaned/disinfected before next resident	O Yes O No	O Yes O No
O Yes O	No	O Yes	O No	O Yes	O No	0 0	Dedicated to resident, cleaned/disinfected before storing Cleaned/disinfected before next resident	O Yes O No	O Yes O No
O Yes O	No	O Yes	O No	O Yes	O No	0 0	Dedicated to resident, cleaned/disinfected before storing Cleaned/disinfected before next resident	O Yes O No	O Yes O No

Notes: ¹Lancet holder devices (e.g., lancing penlets) are not suitable for multi-patient use.

² If the manufacturer does not provide instructions for cleaning and disinfection, then the testing meter should not be used for >1 patient.

³Gloves should be changed and HH performed before assisting the next resident with POCT.